Start with Fluomizin®

10mg Dequalinium Chloride

Reduce pathogenic germs in vaginal





Unique, multiple mode of action against all relevant pathogens of vaginal infections ³

Low systemic exposure, good tolerability and high level of safety ³

Can be used in all phases of pregnancy and lactation ^{2 4}

Effective against the biofilm in bacterial vaginal infections ¹

BRIEF PRESCRIBING INFORMATION OF FLUOMIZIN®

INDICATIONS: Vaginal infections of bacterial and mycotic origin (e.g. bacterial vaginosis and candidiasis), Trichomoniasis and Achievement of asepsis before gynaecological operations and deliveries. DOSAGE & METHOD OF ADMINISTRATION: One vaginal tablet daily for six days. The vaginal tablets should be inserted deeply into the vagina in the evenings before retiring. PREGNANCY AND LACTATION: Can be used during pregnancy and lactation. Caution should be exercised when prescribing Fluomizin® to pregnant women in the first trimester. CONTRAINDICATION: Hypersensitivity to the active substance or to any of the excipients. Ulceration of the vaginal epithelium. Young girls who have not reached sexual maturity should not use Fluomizin®. PRECAUTIONS / INTERACTIONS: No interactions are known. ADVERSE EFFECTS: General disorders and administration site conditions. In rare cases, pruritus, burning or redness. However, these adverse reactions can also be associated with symptoms of vaginal infection. Local irritation reactions, such as bleeding surface defects (erosions) in vagina, were reported in individual cases (0.002%). In these cases, the vaginal surface (vaginal epithelium) was predamaged. Treatment should be stopped, and patient should be advised to seek medical attention. Some very rare cases (0.0003%) of fever have been reported. DATE OF REVISION OF PACKAGE INSERT: January 2013.

References

1. Gaspar C, Rolo J, Cerca N, Palmeira-de- Oliveira R, Martinez-de-Oliveira J, Palmeira-de-Oliveira A. Dequalinium Chloride Effectively Disrupts Bacterial Vaginosis (BV) Gardnerella spp. Bio_Ims. Pathogens. 2021 Feb 25;10(3):261. 2. Fluomizin Package Insert January 2013. 3. Mendling W, Weissenbacher ER, Gerber S, Prasauskas V, Grob P. Use of locally delivered dequalinium chloride in the treatment of vaginal infections: a review. Arch.Gynecol. Obstet. 2016;293:469-84. 4. Demina, T.N., O.N. Pilipenko, O.N., Jotenko, B.A., Baksheeva, O.L. The role of anti-microbial therapy in complex treatment of women with miscarriage. 2005.

For Healthcare Professionals only.





DKSH Malaysia Sdn. Bhd. (4476-U) B-11-01, The Ascent, Paradigm, No. 1, Jalan SS 7/26A, Kelana Jaya, 47301 Petaling Jaya, Selangor Darul Ehsan, Malaysia. Phone +60 3 7882 8888, Fax +60 3 7882 5555

